

Realization of the N (odd)-dimensional Quantum Euclidean Space by Differential Operators*

Yun Li[†] and Sicong Jing
*Department of Modern Physics,
 University of Science and Technology of China,
 Hefei, Anhui 230026, P.R.China*

February 1, 2008

Abstract

The quantum Euclidean space $\mathbb{R}_q^N$ is a kind of noncommutative space which is obtained from ordinary Euclidean space $\mathbb{R}^N$ by deformation with parameter q . When N is odd, the structure of this space is similar to $\mathbb{R}_q^3$. Motivated by realization of $\mathbb{R}_q^3$ by differential operators in $\mathbb{R}^3$, we give such realization for $\mathbb{R}_q^5$ and $\mathbb{R}_q^7$ cases and generalize our results to $\mathbb{R}_q^N$ (N odd) in this paper, that is, we show that the algebra of $\mathbb{R}_q^N$ can be realized by differential operators acting on C^∞ functions on undeformed space $\mathbb{R}^N$.

PACS numbers: 02.20.Uw, 02.40.Gh, 02.30.Tb

1 Introduction

Deformation plays an important role in the development of theoretical models. Indeed the passage from Galilei relativity to that of Lorentz as well as transition from classical to quantum mechanics can both be regarded as deformations associated with the dimensional deformation parameter c (the velocity of light) and $\hbar$ (the Plank constant) respectively. With the intensive study of quantum groups and their associated noncommutative geometry, one builds the Euclidean quantum space $\mathbb{R}_q^N$ related to orthogonal quantum group $SO_q(N)$. In Ref.[1] and [2], some physical systems were constructed based on those Euclidean quantum spaces. So with the view of the historical examples in mind one can look on those physical systems as a deformation of ordinary physics with deformation parameter q .

In order to learn more deformed physical system based on the three-dimensional Euclidean quantum space $\mathbb{R}_q^3$, recently the structure of $\mathbb{R}_q^3$ was discussed in

*The project supported by National Natural Science Foundation of China under Grant No. 10075042

[†]Email: yunlee@mail.ustc.edu.cn

detail[3] and in Ref.[4] the coordinate elements of $\mathbb{R}_q^3$ were represented by differential operators in undeformed Euclidean space $\mathbb{R}^3$. This differential realization is not a novel approach. In fact the method of realizing the generators of universal deformed quantum groups in terms of standard differential operators was discussed thoroughly in Ref.[5].

For the general algebra of $\mathbb{R}_q^N$ it was found that the cases N even and N odd are somewhat different. When N is odd the geometrical structure is similar to $\mathbb{R}_q^3$ case. When, on the other hand, N is even, one should add an additional component to the algebra of $\mathbb{R}_q^N$ [6]. Also in Ref.[7] for the sake of the different parity of N the Euclidean quantum space $\mathbb{R}_q^N$ was sorted as B_N (N odd) and D_N (N even) separately and, for B_N case, the commutation relations for coordinates, as the so-called algebra of the Euclidean quantum space $\mathbb{R}_q^N$, are given by

$$\begin{aligned} X_i X_j &= q X_j X_i, \quad -n \leq i < j \leq n, i \neq -j, \\ X_j X_{-j} &= X_{-j} X_j + q\lambda \sum_{k=-n}^{j-1} q^{\rho_j - \rho_k} X_k X_{-k} - \frac{\lambda q^{\rho_j + 1}}{1 + q^{2n-1}} \sum_{k=-n}^n q^{-\rho_k} X_k X_{-k}, \end{aligned} \quad (1)$$

where $n = \frac{N-1}{2}$, $\lambda = q - q^{-1}$ and $\rho_{-i} = i - \frac{1}{2}$, $\rho_0 = 0$, $\rho_i = -i + \frac{1}{2}$ ($i > 0$). The coordinates X_i satisfy the following conjugation relation

$$\overline{X_i} = q^{\rho_{-i}} X_{-i}. \quad (2)$$

In this paper we restrict our discussion on such B_N case and propose to give a kind of specific realization of the N (odd)-dimensional Euclidean quantum space by differential operators in $\mathbb{R}^N$. We extend the approach developed in Ref.[4] and the result we shall see is somewhat interesting. In section 2 we give a brief review of $N = 3$ case, which is discussed in Ref.[4]. In section 3 we discuss the case of $N = 5$ and 7. In section 4 we will generalize our results to the general N (odd) case.

2 Review of $\mathbb{R}_q^3$ case

If one set $n = 1$ in Eq.(1), one obtains a $SO_q(3)$ -module algebra. This is the algebra of the Euclidean quantum space $\mathbb{R}_q^3$ with definite relations:

$$\begin{aligned} X_{-1} X_0 &= q X_0 X_{-1}, \\ X_0 X_1 &= q X_1 X_0, \\ X_1 X_{-1} &= X_{-1} X_1 + (q^{1/2} - q^{-1/2}) X_0 X_0. \end{aligned} \quad (3)$$

From (2), it is natural to have the conjugation relation of the generators of the above algebra based on $\mathbb{R}_q^3$:

$$\overline{X_1} = q^{1/2} X_{-1}. \quad (4)$$

The next purpose is to represent the algebra in terms of differential operators acting on $\mathbb{R}^3$. For convenience we use polar coordinates (r, θ, φ) and define $\xi = \cos \theta$. An operator that will play a major role in our discussion is:

$$\Lambda_\xi = \frac{1}{2}(\xi \frac{\partial}{\partial \xi} + \frac{\partial}{\partial \xi} \xi) = \xi \frac{\partial}{\partial \xi} + \frac{1}{2} . \quad (5)$$

When acting on L^2 -functions in the common domain of Λ_ξ and Λ_ξ^* ,

$$\Lambda_\xi^* = -\Lambda_\xi \quad (6)$$

will hold. The property of Λ_ξ is followed by

$$\begin{aligned} [\Lambda_\xi, \xi] &= \xi, \\ e^{\alpha \Lambda_\xi} \xi e^{-\alpha \Lambda_\xi} &= e^\alpha \xi . \end{aligned} \quad (7)$$

Now we make an ansatz:

$$\begin{aligned} X_0 &= r\xi, \\ X_1 &= rf(\xi)e^{-\alpha \Lambda_\xi}, \\ X_{-1} &= rg(\xi)e^{\beta \Lambda_\xi} . \end{aligned} \quad (8)$$

Here, $f(\xi)$ and $g(\xi)$ are complex functions acting on $\mathbb{C}^1$. From the first two equations of (3), it is easy to verify

$$e^\alpha = e^\beta = q \quad (9)$$

and from the third equation:

$$f(\xi)g(q^{-1}\xi) - f(q\xi)g(\xi) = (q^{1/2} - q^{-1/2})\xi^2 . \quad (10)$$

If we define

$$\Phi(\xi) = f(\xi)g(q^{-1}\xi), \quad (11)$$

Eq.(10) becomes:

$$\Phi(\xi) - \Phi(q\xi) = (q^{1/2} - q^{-1/2})\xi^2 \quad (12)$$

with the solution:

$$\Phi(\xi) = \Phi(0) - \frac{q^{-1/2}}{1+q}\xi^2 . \quad (13)$$

Next by identifying the radius r with the invariant length in $\mathbb{R}_q^3$:

$$r^2 = R^2 \equiv q^{-1/2}X_{-1}X_1 + X_0X_0 + q^{1/2}X_1X_{-1}, \quad (14)$$

we can determine $\Phi(0)$:

$$\Phi(0) = \frac{q^{1/2}}{1+q} . \quad (15)$$

With the conjugation relation(4), the following equation is obtained

$$g(\xi) = q^{-1/2} \overline{f(q\xi)} \quad (16)$$

and, as a consequence of the definition(11) of $\Phi(\xi)$, we have

$$\Phi(\xi) = q^{-1/2} |f(\xi)|^2 . \quad (17)$$

Combining (13), (15) and (17) leads to the following result

$$\begin{aligned} X_0 &= r\xi, \\ X_1 &= re^{i\varphi} \sqrt{\frac{1}{1+q^{-1}} - \frac{1}{1+q}} \xi^2 q^{-\Lambda_\xi}, \\ X_{-1} &= re^{-i\varphi} \sqrt{\frac{q^{-1}}{1+q^{-1}} - \frac{q}{1+q}} \xi^2 q^{\Lambda_\xi} . \end{aligned} \quad (18)$$

3 $\mathbb{R}_q^5$ and $\mathbb{R}_q^7$ case

When we set $n = 2$ in Eq.(1), we will have the $SO_q(5)$ -module algebra:

$$\begin{aligned} X_i X_j &= q X_j X_i, \quad -2 \leq i < j \leq 2, i \neq -j, \\ X_1 X_{-1} &= X_{-1} X_1 + (q^{1/2} - q^{-1/2}) X_0 X_0, \\ X_2 X_{-2} &= X_{-2} X_2 + q X_1 X_{-1} - q^{-1} X_{-1} X_1, \end{aligned} \quad (19)$$

and the corresponding conjugation relations of the generators are given by

$$\begin{aligned} \overline{X_1} &= q^{1/2} X_{-1}, \\ \overline{X_2} &= q^{3/2} X_{-2} . \end{aligned} \quad (20)$$

To represent the algebra(19) in terms of differential operators acting on the Euclidean space $\mathbb{R}^5$, we also use polar coordinates $(r, \theta_1, \theta_2, \varphi_1, \varphi_2)$ of $\mathbb{R}^5$ and define $\xi_i = \cos \theta_i$ ($i = 1, 2$). Like the ansatz made in Eq.(8) when dealing with $\mathbb{R}_q^3$ case and after considering the explicit expression(18), we assume that the generators X_i have the following form:

$$\begin{aligned}
X_0 &= r\xi_1, \\
X_1 &= re^{i\varphi_1}f_1(\xi_1, \xi_2)e^{-\alpha_1\Lambda_{\xi_1}}, \\
X_{-1} &= re^{-i\varphi_1}g_1(\xi_1, \xi_2)e^{\beta_1\Lambda_{\xi_1}}, \\
X_2 &= re^{i\varphi_2}f_2(\xi_1, \xi_2)e^{-\alpha_2(\Lambda_{\xi_1}+\Lambda_{\xi_2})}, \\
X_{-2} &= re^{-i\varphi_2}g_2(\xi_1, \xi_2)e^{\beta_2(\Lambda_{\xi_1}+\Lambda_{\xi_2})}.
\end{aligned} \tag{21}$$

Here, $f_{1,2}(\xi_1, \xi_2)$ and $g_{1,2}(\xi_1, \xi_2)$ are absolutely real functions acting on $\mathbb{R}^2$ because we write the complex parts into exponential forms in the above expressions. From the first equation of (19), the following

$$e^{\alpha_i} = e^{\beta_i} = q, \quad (i = 1, 2) \tag{22}$$

also holds. Simultaneously some confining conditions related to the functions $f_{1,2}(\xi_1, \xi_2)$ and $g_{1,2}(\xi_1, \xi_2)$ are obtained. We list them below

$$\begin{aligned}
g_1(q\xi_1, q\xi_2)g_2(\xi_1, \xi_2) &= qg_1(\xi_1, \xi_2)g_2(q\xi_1, \xi_2), \\
f_1(q\xi_1, q\xi_2)g_2(\xi_1, \xi_2) &= qf_1(\xi_1, \xi_2)g_2(q^{-1}\xi_1, \xi_2), \\
g_1(\xi_1, \xi_2)f_2(q\xi_1, \xi_2) &= qg_1(q^{-1}\xi_1, q^{-1}\xi_2)f_2(\xi_1, \xi_2), \\
f_1(\xi_1, \xi_2)f_2(q^{-1}\xi_1, \xi_2) &= qf_1(q^{-1}\xi_1, q^{-1}\xi_2)f_2(\xi_1, \xi_2).
\end{aligned} \tag{23}$$

From the second equation of (19), we have

$$f_1(\xi_1, \xi_2)g_1(q^{-1}\xi_1, \xi_2) - f_1(q\xi_1, \xi_2)g_1(\xi_1, \xi_2) = (q^{1/2} - q^{-1/2})\xi_1^2. \tag{24}$$

With the definition:

$$\Phi_1(\xi_1, \xi_2) = f_1(\xi_1, \xi_2)g_1(q^{-1}\xi_1, \xi_2), \tag{25}$$

this equation becomes:

$$\Phi_1(\xi_1, \xi_2) - \Phi_1(q\xi_1, \xi_2) = (q^{1/2} - q^{-1/2})\xi_1^2 \tag{26}$$

and has the solution:

$$\Phi_1(\xi_1, \xi_2) = \Phi_1(0, 0) - \frac{q^{-1/2}}{1+q}\xi_1^2 + B_1\xi_2^2, \tag{27}$$

in which $\Phi_1(0, 0)$ and B_1 need to be determined further. The third equation of (19) leads to the results:

$$\begin{aligned}
&f_2(\xi_1, \xi_2)g_2(q^{-1}\xi_1, q^{-1}\xi_2) \\
&= f_2(q\xi_1, q\xi_2)g_2(\xi_1, \xi_2) + qf_1(\xi_1, \xi_2)g_1(q^{-1}\xi_1, \xi_2) - q^{-1}f_1(q\xi_1, \xi_2)g_1(\xi_1, \xi_2).
\end{aligned} \tag{28}$$

By defining

$$\Phi_2(\xi_1, \xi_2) = f_2(\xi_1, \xi_2)g_2(q^{-1}\xi_1, q^{-1}\xi_2), \quad (29)$$

we have the following

$$\Phi_2(\xi_1, \xi_2) = \Phi_2(q\xi_1, q\xi_2) + q\Phi_1(\xi_1, \xi_2) - q^{-1}\Phi_1(q\xi_1, \xi_2). \quad (30)$$

Substituting the solution(27) of $\Phi_1(\xi_1, \xi_2)$ into the above equation, consequently we have the solution of $\Phi_2(\xi_1, \xi_2)$:

$$\Phi_2(\xi_1, \xi_2) = \Phi_2(0, 0) - q^{-1}B_1\xi_2^2 \quad (31)$$

and also we can determine the value of $\Phi_1(0, 0)$ satisfying

$$\Phi_1(0, 0) = 0. \quad (32)$$

Next also by identifying radius r with the invariant length in $\mathbb{R}_q^5$:

$$r^2 = R^2 \equiv q^{-3/2}X_{-2}X_2 + q^{-1/2}X_{-1}X_1 + X_0X_0 + q^{1/2}X_1X_{-1} + q^{3/2}X_2X_{-2} \quad (33)$$

and after a careful calculation, $\Phi_2(0, 0)$ can be determined:

$$\Phi_2(0, 0) = \frac{1}{q^{3/2} + q^{-3/2}}. \quad (34)$$

So we have

$$\begin{aligned} \Phi_1(\xi_1, \xi_2) &= B_1\xi_2^2 - \frac{q^{-1/2}}{1+q}\xi_1^2, \\ \Phi_2(\xi_1, \xi_2) &= \frac{1}{q^{3/2} + q^{-3/2}} - q^{-1}B_1\xi_2^2 \end{aligned} \quad (35)$$

only with parameter B_1 needing to be determined further. By applying the conjugation relations(20), we derive the following equations

$$\begin{aligned} \Phi_1(\xi_1, \xi_2) &= q^{-1/2}f_1^2(\xi_1, \xi_2), \\ \Phi_2(\xi_1, \xi_2) &= q^{-3/2}f_2^2(\xi_1, \xi_2). \end{aligned} \quad (36)$$

With (35) and (36) it is easy to obtain

$$\begin{aligned}
X_0 &= r\xi_1, \\
X_1 &= re^{i\varphi_1} \sqrt{q^{1/2}B_1\xi_2^2 - \frac{1}{1+q}\xi_1^2 q^{-\Lambda_{\xi_1}}}, \\
X_{-1} &= re^{-i\varphi_1} \sqrt{q^{-1/2}B_1\xi_2^2 - \frac{q}{1+q}\xi_1^2 q^{\Lambda_{\xi_1}}}, \\
X_2 &= re^{i\varphi_2} \sqrt{\frac{1}{1+q^{-3}} - q^{1/2}B_1\xi_2^2 q^{-(\Lambda_{\xi_1}+\Lambda_{\xi_2})}}, \\
X_{-2} &= re^{-i\varphi_2} \sqrt{\frac{q^{-3}}{1+q^{-3}} - q^{-1/2}B_1\xi_2^2 q^{\Lambda_{\xi_1}+\Lambda_{\xi_2}}}.
\end{aligned} \tag{37}$$

By setting

$$B_1 = \frac{q^{-1/2}}{1+q^3}, \tag{38}$$

the above expression can have a uniform expression with (18). So the generators X_i of the $SO_q(5)$ -module algebra can be written as

$$\begin{aligned}
X_0 &= r\xi_1, \\
X_1 &= re^{i\varphi_1} \sqrt{\frac{1}{1+q^3}\xi_2^2 - \frac{1}{1+q}\xi_1^2 q^{-\Lambda_{\xi_1}}}, \\
X_{-1} &= re^{-i\varphi_1} \sqrt{\frac{q^{-1}}{1+q^3}\xi_2^2 - \frac{q}{1+q}\xi_1^2 q^{\Lambda_{\xi_1}}}, \\
X_2 &= re^{i\varphi_2} \sqrt{\frac{1}{1+q^{-3}} - \frac{1}{1+q^3}\xi_2^2 q^{-(\Lambda_{\xi_1}+\Lambda_{\xi_2})}}, \\
X_{-2} &= re^{-i\varphi_2} \sqrt{\frac{q^{-3}}{1+q^{-3}} - \frac{q^{-1}}{1+q^3}\xi_2^2 q^{\Lambda_{\xi_1}+\Lambda_{\xi_2}}}.
\end{aligned} \tag{39}$$

The above expressions are obviously consistent with the confining conditions metioned in Eq.(23). Our purpose is to represent the algebra of the N (odd)-dimensional Euclidean quantum space in terms of differential operators acting on $\mathbb{R}^N$, so if we obtain the representation in $\mathbb{R}_q^7$ case again, we can find the intrinsic rule of expression forms and generalize our results to the N (odd) case. For $N = 7$ case, the algebra relations are given by

$$\begin{aligned}
X_i X_j &= q X_i X_j, \quad -3 \leq i < j \leq 3, i \neq -j, \\
X_1 X_{-1} &= X_{-1} X_1 + (q^{1/2} - q^{-1/2}) X_0 X_0, \\
X_2 X_{-2} &= X_{-2} X_2 + q X_1 X_{-1} - q^{-1} X_{-1} X_1, \\
X_3 X_{-3} &= X_{-3} X_3 + q X_2 X_{-2} - q^{-1} X_{-2} X_2
\end{aligned} \tag{40}$$

with the conjugation relations of the generators:

$$\begin{aligned}\overline{X}_1 &= q^{1/2} X_{-1}, \\ \overline{X}_2 &= q^{3/2} X_{-2}, \\ \overline{X}_3 &= q^{5/2} X_{-3}.\end{aligned}\tag{41}$$

Following the same procedure as we obtain expression(39), we have the following

$$\begin{aligned}X_0 &= r\xi_1, \\ X_1 &= re^{i\varphi_1} \sqrt{\frac{1}{1+q^3}\xi_2^2 - \frac{1}{1+q}\xi_1^2 q^{-\Lambda_{\xi_1}}}, \\ X_{-1} &= re^{-i\varphi_1} \sqrt{\frac{q^{-1}}{1+q^3}\xi_2^2 - \frac{q}{1+q}\xi_1^2 q^{\Lambda_{\xi_1}}}, \\ X_2 &= re^{i\varphi_2} \sqrt{\frac{1}{1+q^5}\xi_3^2 - \frac{1}{1+q^3}\xi_2^2 q^{-(\Lambda_{\xi_1}+\Lambda_{\xi_2})}}, \\ X_{-2} &= re^{-i\varphi_2} \sqrt{\frac{q^{-3}}{1+q^5}\xi_3^2 - \frac{q^{-1}}{1+q^3}\xi_2^2 q^{\Lambda_{\xi_1}+\Lambda_{\xi_2}}}, \\ X_3 &= re^{i\varphi_3} \sqrt{\frac{1}{1+q^{-5}} - \frac{1}{1+q^5}\xi_3^2 q^{-(\Lambda_{\xi_1}+\Lambda_{\xi_2}+\Lambda_{\xi_3})}}, \\ X_{-3} &= re^{-i\varphi_3} \sqrt{\frac{q^{-5}}{1+q^{-5}} - \frac{q^{-3}}{1+q^5}\xi_3^2 q^{\Lambda_{\xi_1}+\Lambda_{\xi_2}+\Lambda_{\xi_3}}}.\end{aligned}\tag{42}$$

Also here $(r, \theta_1, \theta_2, \theta_3, \varphi_1, \varphi_2, \varphi_3)$ represent polar coordinates of the Euclidean space $\mathbb{R}^7$ and the definitions of ξ_i and Λ_{ξ_i} are same as (18) and (39).

4 $\mathbb{R}_q^N$ (N odd) case

Now we deal with the algebra of $\mathbb{R}_q^N$ (N odd) and give the differential realization of it. After speculating the realization in $N = 3, 5$ and 7 cases, it is not difficult to find that for the general N (odd) case the realization can be expressed as

$$\begin{aligned}
X_0 &= r\xi_1, \\
X_i &= re^{i\varphi_i} \sqrt{\frac{1}{1+q^{2i+1}}\xi_{i+1}^2 - \frac{1}{1+q^{2i-1}}\xi_i^2} q^{-\sum_{m=1}^i \Lambda_{\xi_m}}, \\
X_{-i} &= re^{-i\varphi_i} \sqrt{\frac{q^{-2i+1}}{1+q^{2i+1}}\xi_{i+1}^2 - \frac{q^{-2i+3}}{1+q^{2i-1}}\xi_i^2} q^{\sum_{m=1}^i \Lambda_{\xi_m}}, \quad 1 \leq i \leq n-1, n = \frac{N-1}{2}, \\
X_n &= re^{i\varphi_n} \sqrt{\frac{1}{1+q^{-2n+1}} - \frac{1}{1+q^{2n-1}}\xi_n^2} q^{-\sum_{m=1}^n \Lambda_{\xi_m}}, \\
X_{-n} &= re^{-i\varphi_n} \sqrt{\frac{q^{-2n+1}}{1+q^{-2n+1}} - \frac{q^{-2n+3}}{1+q^{2n-1}}\xi_n^2} q^{\sum_{m=1}^n \Lambda_{\xi_m}}.
\end{aligned} \tag{43}$$

We also assume that $(r, \theta_1, \dots, \theta_i, \dots, \theta_n, \varphi_1, \dots, \varphi_i, \dots, \varphi_n)$ are polar coordinates of the Euclidean space $\mathbb{R}^N$ and have $\xi_i = \cos \theta_i, \Lambda_{\xi} = \xi \frac{\partial}{\partial \xi} + \frac{1}{2}$ in the above expression.

Next we will verify that the above expressions are definitely a kind of differential realization for the general N (odd) case. One can rewrite the last line of the commutative relations(1) in a more simple form[8]:

$$X_s X_{-s} = X_{-s} X_s + \lambda q^{s-3/2} \left(\sum_{i=1}^{s-1} q^{\rho_i} X_{-i} X_i + \frac{q}{1+q} X_0 X_0 \right), \quad s > 0 \tag{44}$$

For $s = 1$, it becomes

$$X_1 X_{-1} = X_{-1} X_1 + (q^{1/2} - q^{-1/2}) X_0 X_0, \tag{45}$$

For $s > 1$, this relation is equivalent to the following one

$$X_s X_{-s} = X_{-s} X_s + q X_{s-1} X_{-s+1} - q^{-1} X_{-s+1} X_{s-1} \tag{46}$$

So we can rewrite the algebra of $\mathbb{R}_q^N$ (N odd)(1) as following

$$\begin{aligned}
X_i X_j &= q X_j X_i, \quad -n \leq i < j \leq n, i \neq -j, \\
X_1 X_{-1} &= X_{-1} X_1 + (q^{1/2} - q^{-1/2}) X_0 X_0, \\
X_s X_{-s} &= X_{-s} X_s + q X_{s-1} X_{-s+1} - q^{-1} X_{-s+1} X_{s-1}, \quad 1 < s \leq n
\end{aligned} \tag{47}$$

Now substituting the expressions(43) into the above relations we will find that the right-hand side of each line in the relations is equal to the left-hand one. So Eq.(43) is a possible realization by standard differential operators in $\mathbb{R}^N$. To make sense, we take the limit $q \rightarrow 1$ in (43) and find that X_i ($-n \leq i \leq n$) become linearly independent. In this limit the deformed quantum space $\mathbb{R}_q^N$ reduces into the undeformed $\mathbb{R}^N$ and the generators X_i could be a set of basis of $\mathbb{R}^N$.

References

- [1] U. Carow-Watamura, M. Schliker and S. Watamura, Z. Phys. C **49**, 439 (1991)
- [2] H. Ocampo, Z.Phys.C **70**, 525 (1996)
- [3] B. L. Cerchiai, J. Madore, S. Schraml and J. Wess, Eur. Phys. J. C **16**, 169 (2000)
- [4] S. Schraml and J. Wess, Eur. Phys. J. C **17**, 353 (2000)
- [5] T. L. Curtright and C. K. Zachos, Phys. Lett B **243**, 237 (1990)
- [6] B. L. Cerchiai, G. Fiore and J. Madore, Commun. Math. Phys. **217**, 521 (2001)
- [7] L. D. Faddev, N. Y. Reshetikhin and L. A. Takhtajan, Algebra Anal. **1**, 178 (1989)
- [8] O. Ogievetsky, Lett. Math. Phys. **24**, 245 (1992)